


Important for Preventing the Spread of the Virus

Examples of Best Practices, a "New Lifestyle"

HAMAMATSU CITY



**Keep
a 2m distance**



Wear masks



Wash hands



Meet friends outdoors



**Avoid face to face
conversations**



出世大名
家康くん
出世法師
直虎ちゃん

Lifestyle
(e.g.)

Shopping



**Go alone
or in a small group
at less crowded times**

Exercise



Using videos at home

Eating Out



**Takeout
or order deliveries**

Work Life
(e.g.)



Work from home



**Stagger
your commute times**



Hold meetings online

5 Situations with a High Risk of Infection

① Social Events with Alcohol

- After drinking alcohol, many people become excited and their attentiveness decreases. In addition, their hearing is dulled so they are more likely to speak loudly.
- The risk of infection increases especially when large numbers of people stay in small partitioned spaces for long periods of time.
- The risk of infection is also high if you pass drinks around and share chopsticks.



② Eating and Drinking with Large Groups of People or for Long Periods of Time

- The risk of infection is higher if you have a meal over a long period of time, or while entertaining guests or go barhopping as compared to a short meal.
- The risk also increases if you have a meal with a large group of people (more than 4 people) because droplets may spread due to speaking loudly.



③ Talking Without a Mask

- The risk of infection via water droplets increases when you talk in close distances without wearing masks.
- There have been cases of infection caused by singing karaoke without a mask on during the day.
- Please be careful even on buses or cars.



④ Living Together in Small Spaces

- There is an increased risk of infection if you live with other people in small spaces and spend long periods of time together in enclosed spaces.
- There have been cases of infection that are suspected to have arisen from shared spaces like dormitories or toilets.



⑤ Change in Locations

- There may be a higher risk of infection when you change locations during break time due to lowering your guard and a change in surroundings.
- Cases of infection arising from spaces like the break room, smoking area or changing room have been confirmed.



Complete Measures to Prevent Infection during Events and Dinner Parties

Please refrain from participating in events and dinner parties which fall under the any of the 3Cs (closed space, crowds and close contact) or fail to be thorough in setting up basic preventive measures.

In particular, please refrain from participating in events and parties with large crowds of people with people shouting loudly.

If you will be participating in such events or parties, please take appropriate measures to prevent infection by securing a safe distance from other people, sanitizing your hands, wearing a mask and refraining from speaking in a loud manner.

Please consider options such as spending time at home with your family or participating in online activities instead.

Please refrain from heavy drinking on the street and in shops or participating in drinking parties.